



TECHNICAL DATA SHEET



Gentamicine sulfate

AGRICULTURAL BACTERICIDE

SOLUBLE CONCENTRATE (SL)

Agry Gent Plus® LC is the new liquid bactericide with systemic action and broad spectrum of the AGRY GENT's family, its active ingredient is Gentamicin Sulfate contained in a formulation focused on the prevention and control of vascular bacteria in the soil.

The product can be applied in any irrigation systems.

PERCENTAGE COMPOSITION

Active ingredients:

Gentamicine sulfate8%

Inert ingredients:

Diluent and dispersants92.0%

ADVANTAGES:

- ✓ Prevents and controls bacterial diseases
- ✓ Enfoque innovador en el suelo
- ✓ Easy to dose
- ✓ Safe to use
- ✓ Stable formulation
- ✓ Easier application in pressurized irrigation systems due to its formulation without inert solids
- ✓ Ideal for dipping treatments of bulbs, stem cuttings and tubers

PACKING

Agry Gent Plus® LC presentation: 1 L, soluble concentrate.

PRODUCT PREPARATION

Pour the product into the tank with half the recommended water, then add the other half of the water and stir the mixture constantly.

The product can be applied in drench directly to the base of the plants, or sprayed directly to the soil at the time of sowing or another vegetative stage and through any irrigation system.

Apply it immediately after preparing the mixture.





AUTHORIZED DOSE AND CROPS:

AUTHORIZED CROPS	DISEASE	DOSE	REMARKS
Pear (21)	Fire blight <i>Erwinia amylovora</i>	300 mg/5ml de agua equivalente a 5ml / árbol	Application by injection to the tree trunk
Summer squash, pumpkin, chayote, melon, cucumber y watermelon	Angular leaf spot <i>Pseudomonas syringae pv. lachrymans</i>	150 - 500 mL/ ha	Make three in-drench applications at 7-day intervals when the first symptoms of the disease are detected or when environmental conditions are favorable for its development, suggested application volume 1000 L of water/ha.
Eggplant, pepper, bell pepper, tomato, green tomato y potato (SL)	Bacterial wilt <i>Ralstonia solanacearum</i>	150 - 500 mL/ ha	Make three in-drench applications at 7-day intervals when the first symptoms of the disease are detected or when environmental conditions are favorable for its development, suggested application volume 1000 L of water/ha.

() PHI. Preharvest Interval. SL = Without limit

CONSIDERATIONS

Do not apply **Agry Gent Plus® LC** with wind speeds greater than 15 km/hour.

COMPATIBILITY

Do not mix **Agry Gent Plus® LC** with Bordeaux Broth and other products with a strong alkaline reaction. When mixtures are required to be made, they can only be made with products that have a current registration and are authorized for the indicated crops here.

PHYTOTOXICITY

Agry Gent Plus® LC is not phytotoxic to the indicated crops here, if applied according to label recommendations.

RESISTANCE MANAGEMENT

“TO PREVENT THE DEVELOPMENT OF RESISTANT POPULATIONS, ALWAYS RESPECT THE DOSE AND FREQUENCY OF APPLICATION; AVOID THE REPEATED USE OF THIS PRODUCT, ALTERNATING IT WITH OTHER CHEMICAL GROUPS WITH DIFFERENT MODES OF ACTION AND DIFFERENT DETOXIFICATION MECHANISMS AND THROUGH THE SUPPORT OF OTHER CONTROL METHODS”

MANUFACTURER AND REGISTRY HOLDER

Química Agronómica de México, S. de R.L. de C.V.
Calle 18 No. 20501, Parque Industrial Impulso, C.P. 31183
Chihuahua, Chihuahua, México
Tel. +52 614 483 9000

